arXiv:0810.0511v2 [hep-ph] 16 Oct 2008

Lightest Neutralino Mass in the MNSSM

S. Hesselbach, G. Moortgat-Pick
IPPP, Durham, DH1 3LE, UK
D. J. Miller, R. Nevzorov
University of Glasgow, Glasgow, G12 8QQ, UK
M. Trusov
ITEP, Moscow, 117218, Russia

We argue that the allowed range of the mass of the lightest neutralino in the MNSSM is limited. We establish the theoretical upper bound on the lightest neutralino mass and obtain an approximate solution for this mass.

1. INTRODUCTION

Recent observations indicate that $22\% - 25\%$ of the energy density of the Universe exists in the form of stable non–baryonic, non–luminos (dark) matter. Supersymmetric (SUSY) models provide a good candidate for the cold dark matter component of the Universe. If R–parity is conserved, the lightest neutralino is absolutely stable and can play the role of dark matter. However the minimal supersymmetric standard model (MSSM) suffers from the so-called μ–problem. The MSSM superpotential contains only one bilinear term $\mu(\hat{H}_1\epsilon\hat{H}_2)$. The parameter μ is expected to be of the order of M_{Pl} or GUT scale. At the same time the correct pattern of electroweak (EW) symmetry breaking requires $\mu \sim 100 - 1000\,\mathrm{GeV}$. The μ–problem can be solved within the minimal non–minimal supersymmetric standard model (MNSSM). The superpotential of the MNSSM can be written as

$$W = \lambda\hat{S}(\hat{H}_d\epsilon\hat{H}_u) + \xi\hat{S} + W_{MSSM}(\mu = 0)\,. \tag{1}$$

At the EW scale the singlet superfield $\hat{S}$ gets non-zero vacuum expectation value (VEV) $\langle S\rangle = s/\sqrt{2}$ and an effective μ-term ($\mu_{eff} = \lambda s/\sqrt{2}$) is generated. Thus in the MNSSM the μ problem is solved in a similar way to the next–to–minimal supersymmetric standard model (NMSSM), but without the accompanying problems of singlet tadpoles or domain walls.

2. UPPER BOUND ON THE LIGHTEST NEUTRALINO MASS

The neutralino sector of the MNSSM is formed by the superpartners of the neutral gauge bosons ($\tilde{W}_3$, $\tilde{B}$) and neutral Higgsino fields ($\tilde{H}^0_d, \tilde{H}^0_u, \tilde{S}$). In the field basis ($\tilde{B}$, $\tilde{W}_3$, $\tilde{H}^0_d$, $\tilde{H}^0_u$, $\tilde{S}$) the neutralino mass matrix takes a form

$$M_{\tilde{\chi}} = \begin{pmatrix} M_1 & 0 & -M_Z s_W c_\beta & M_Z s_W s_\beta & 0 \\ 0 & M_2 & M_Z c_W c_\beta & -M_Z c_W s_\beta & 0 \\ -M_Z s_W c_\beta & M_Z c_W c_\beta & 0 & -\mu_{eff} & -\frac{\lambda v}{\sqrt{2}} s_\beta \\ M_Z s_W s_\beta & -M_Z c_W s_\beta & -\mu_{eff} & 0 & -\frac{\lambda v}{\sqrt{2}} c_\beta \\ 0 & 0 & -\frac{\lambda v}{\sqrt{2}} s_\beta & -\frac{\lambda v}{\sqrt{2}} c_\beta & 0 \end{pmatrix}, \tag{2}$$

where $s_W = \sin\theta_W$, $c_W = \cos\theta_W$, $s_\beta = \sin\beta$, and $c_\beta = \cos\beta$. Here $\tan\beta = v_2/v_1$ and $v = \sqrt{v_1^2 + v_2^2} = 246\,\mathrm{GeV}$, where v_1 and v_2 are the VEVs of H_d and H_u, respectively. The neutralino spectrum in the MNSSM may be parametrised in terms of

$$\lambda\,, \qquad \mu_{eff}\,, \qquad \tan\beta\,, \qquad M_1\,, \qquad M_2\,. \tag{3}$$

In the minimal SUGRA inspired SUSY GUT's $M_1 \simeq 0.5M_2$. The validity of perturbation theory up to the GUT scale implies that $\lambda(M_Z) \le 0.7$. When λ is small the non–observation of Higgs bosons at LEP rules out $\tan\beta \le 2.5$. LEP searches for SUSY particles also set lower bounds on $|M_2|$, $|\mu_{eff}| > 90-100\,\mathrm{GeV}$.

In order to find theoretical bounds on the neutralino masses $m_{\chi_i^0}$ it is convenient to consider the matrix $M_{\tilde{\chi}^0}M^{\dagger}_{\tilde{\chi}^0}$. In the basis $\left(\tilde{B}, \tilde{W}_3, -\tilde{H}_d^0 s_\beta + \tilde{H}_u^0 c_\beta, \tilde{H}_d^0 c_\beta + \tilde{H}_u^0 s_\beta, \tilde{S}\right)$ the bottom-right 2×2 block of $M_{\tilde{\chi}^0}M^{\dagger}_{\tilde{\chi}^0}$ takes the form

$$\begin{pmatrix} |\mu_{eff}|^2 + \sigma^2 & \nu^*\mu_{eff} \\ \nu\mu^*_{eff} & |\nu|^2 \end{pmatrix}, \tag{4}$$

where $\sigma^2 = M_Z^2\cos^2 2\beta + |\nu|^2\sin^2 2\beta$, $\nu = \lambda v/\sqrt{2}$. Since the minimal eigenvalue of any hermitian matrix is less than its smallest diagonal element at least one neutralino in the MNSSM is always light, i.e. $|m_{\chi_1^0}| \le |\nu|$. Therefore in contrast with the MSSM the lightest neutralino in the MNSSM remains light even when the SUSY breaking scale tends to infinity. The mass of the lightest neutralino must be also smaller than the minimal eigenvalue μ_0^2 of the submatrix (4), i.e.

$$|m_{\chi_1^0}|^2 \le \mu_0^2 = \frac{1}{2}\left[|\mu_{eff}|^2 + \sigma^2 + |\nu|^2 - \sqrt{\left(|\mu_{eff}|^2 + \sigma^2 + |\nu|^2\right)^2 - 4|\nu|^2\sigma^2}\right]. \tag{5}$$

The value of μ_0 decreases with increasing $|\mu_{eff}|$. Taking into account the restrictions on $|\mu_{eff}|$ and $\lambda(M_Z)$ we find that $|m_{\chi_1^0}| \le 80-85\,\mathrm{GeV}$ [1]. The obtained theoretical restriction on $|m_{\chi_1^0}|$ allows to discriminate the MNSSM from other SUSY models.

3. APPROXIMATE SOLUTION

The mass of the lightest neutralino can be computed numerically by solving the characteristic equation

$$\begin{aligned} \det\left(M_{\tilde{\chi}^0} - \kappa I\right) = \Big(M_1M_2 - (M_1+M_2)\kappa + \kappa^2\Big)\Big(\kappa^3 - (\mu_{eff}^2 + \nu^2)\kappa + \nu^2\mu_{eff}\sin 2\beta\Big) \\ + M_Z^2\Big(\tilde{M} - \kappa\Big)\Big(\kappa^2 + \mu_{eff}\sin 2\beta\kappa - \nu^2\Big) = 0\,, \end{aligned} \tag{6}$$

where $\tilde{M} = M_1 c_W^2 + M_2 s_W^2$. Because in the MNSSM $|m_{\chi_1^0}|$ is normally much smaller than the masses of the other neutralinos one can ignore κ^3, κ^4 and κ^5 terms in the characteristic equation so that it reduces to

$$\kappa^2 - B\kappa + C = 0, \tag{7}$$

where

$$B = \frac{M_1M_2}{M_1+M_2} + \left(\frac{\nu^2}{\mu_{eff}^2+\nu^2} - \frac{M_Z^2}{\mu_{eff}^2+\nu^2}\frac{\tilde{M}}{M_1+M_2}\right)\mu_{eff}\sin 2\beta - \frac{M_Z^2\nu^2}{(M_1+M_2)(\mu_{eff}^2+\nu^2)}\,,$$

$$C = \frac{\nu^2}{\mu_{eff}^2+\nu^2}\left(\frac{M_1M_2}{M_1+M_2}\mu_{eff}\sin 2\beta - \frac{\tilde{M}}{M_1+M_2}M_Z^2\right).$$

Then the mass of the lightest neutralino can be approximated by [1]

$$|m_{\chi_1^0}| = \mathrm{Min}\left\{\frac{1}{2}\left|B - \sqrt{B^2-4C}\right|, \frac{1}{2}\left|B + \sqrt{B^2-4C}\right|\right\}. \tag{8}$$

As follows from Fig. 1–2 the approximate solution describes the numerical one with relatively high accuracy. With increasing μ_{eff} and $M_{1,2}$ the lightest neutralino mass decreases. If either $|\mu_{eff}|$ or $M_{1,2} \gg M_Z$ then

$$|m_{\chi_1^0}| \simeq \frac{|\mu_{eff}|\nu^2\sin 2\beta}{\mu_{eff}^2+\nu^2}\,. \tag{9}$$

According to Eq.(9) $m_{\chi_1^0}$ is inversely proportional to μ_{eff}. It decreases when $\tan\beta$ grows and vanishes when λ tends to zero. When $\lambda \to 0$ the mass of the lightest neutralino is proportional to λ^2. When $|m_{\chi_1^0}| \ll M_Z$ the lightest neutralino is predominantly singlino that makes its direct observation at future colliders rather challenging.



Figure 1: Lightest neutralino mass versus μ_{eff} for $\lambda = 0.7$, $M_1 = 0.5M_2$, $M_2 = 200\,\mathrm{GeV}$. Solid and dashed lines correspond to $\tan\beta = 10$ and 3. Upper and lower dashed–dotted lines represent the theoretical restriction, Eq. (5), on $|m_{\chi_1^0}|$ for $\tan\beta = 3$ and 10. Dotted lines correspond to the approximate solution, Eq. (8).



Figure 2: Lightest neutralino mass as a function of M_2 for $\lambda = 0.7$, $M_1 = 0.5M_2$, $\mu_{eff} = 200\,\mathrm{GeV}$. The notations are the same as in Fig. 1.

Acknowledgments

RN would like to thank A. Pilaftsis and M. Vysotsky for fruitful discussions.
RN acknowledges support from the SHEFC grant HR03020 SUPA 36878.

References

[1] S. Hesselbach, D.J. Miller, G. Moortgat-Pick, R. Nevzorov, M. Trusov, "Theoretical upper bound on the mass of the LSP in the MNSSM", Phys. Lett. B **662** (2008) 199.